

Short Commentary

Open Access, Volume 2

Pros and cons of iron deficiency and iron deficiency anemia

*Corresponding Author: Adel Ekladiou

Faculty of Health and Medical Sciences University of
Western Australia 35 Stirling Hwy, Crawley, Western
Australia.

Royal Hobart Hospital, 48 Liverpool Street, Tasmania
7000, Australia.

Tel: 61-499-449-905; Email: ekladiou@gmail.com

Introduction

Iron deficiency anemia (IDA) and iron deficiency (AD) are very common specially in low-income countries [1]. People can suffer from symptoms of anemia in the absence of anemia itself if they have decreased iron stores as red blood cells and tissue will not receive enough iron from the plasma when iron stores are low.

Received: Nov 29, 2021

Accepted: Jan 06, 2022

Published: Jan 13, 2022

Archived: www.jjgastro.com

Copyright: © Ekladiou A (2022).

Discussion

During inflammation, the hepcidin level increases and ferroprotein transcription decreases which results in decreased recycling of iron to the plasma and iron absorption from the duodenum [2]. Iron is essential for physiological and cellular functions including formation and repair of DNA, enzymatic activity, mitochondrial function and neurotransmission [3]. Iron deficiency should be recognized early, investigated, and treated to improve quality of life in many patient groups including patients with chronic heart failure, chronic kidney disease, inflammatory bowel disease, and pregnant women in the second and third trimester [3]. Common debilitating symptoms due to ID include fatigue, tiredness, lethargy, dizziness, tinnitus, headache, pallor, difficulty concentrating and restless leg syndrome which can be treated by iron supplementation even in the absence of anemia [4]. Preoperative anemia is a predictor of high mortality and morbidity even in the absence of symptoms due to IDA [5]. Body iron in red cells is 2.5 grams, 130 mg in myoglobin and 150 mg in enzymes [4]. Average iron stores in men are 9.7 mg/kg, 5.7 mg/kg in premenopausal women and 7.8 mg/kg in post-menopausal women in high income countries [5]. One per cent of the total body iron is bound in the plasma to transferrin

which can bind to the tissue receptor called soluble transferrin receptor. Plasma iron is rapidly recycled from aged red blood cells by macrophages [6]. 1-2 mg of daily iron is absorbed from the duodenum.

Hepcidin is a liver enzyme which is essential for regulation of systemic iron hemostasis. Iron is regularly exported to the plasma from macrophages, the duodenum and the liver [6]. During iron deficiency, the level of hepcidin drops leading to increased iron absorption and increased recycling. During inflammation, increased levels of hepcidin lead to decreased iron absorption from the duodenum and decreased recycling of iron leading to a functional deficiency, as iron stores are not affected. In iron deficiency, hepcidin reduces absorption of iron, which subsequently renders oral iron ineffective. As such, patients should be treated with a parenteral infusion [7,8].

One milliliter of blood contains 0.5 mg of iron. For each donation of blood, a person loses 250 mg of iron. As such, frequent donors should receive iron transfusions after donating blood [9]. The average daily intake of iron is 8 mg in adult males, 18 mg in premenopausal women and 27 mg in pregnant women [10]. Haem iron is more efficient than non-haem iron [10]. Vitamin C increases the absorption of the non-haem iron, and tan-

nic acid (in tea and coffee) decreases absorption of iron. Gastric acidity increases the need for iron absorption. Proton pump inhibitors and H₂ blockers adversely affect oral absorption of iron. *Helicobacter pylori* decreases absorption of iron and causes iron deficiency anemia. Eradication of *H Pylori* subsequently cures iron deficiency and iron deficiency anemia [10,11]. Bariatric surgery, whether gastric bypass or gastric sleeve, is a common cause of ID and IDA [12]. Obese people develop functional iron deficiency due to inflammation, leading to nonalcoholic steatohepatitis, subsequently increasing the ferritin. This increases the hepcidin, leading to impaired iron absorption and recycling. Premenopausal women should not be offered endoscopy unless they do not improve after iron supplementation, or if they have a family history of colorectal cancer in the first degree relative and investigations for coeliac disease are negative.

Patients can develop iron deficiency in coeliac disease even with normal endoscopy, and usually histology shows villous atrophy and crypt hyperplasia. Transglutaminase serum IgA antibody and serum IgA are the standard of care serology. Endomysial antibodies are very specific but less sensitive for diagnosing coeliac disease. Males over fifty years and postmenopausal women should have upper and lower endoscopy to exclude gastric ulceration, autoimmune gastritis, polyp, colorectal cancer, diverticulosis, and angiodysplasia. People from areas endemic for *Plasmodium falciparum* should receive prophylaxis treatment for malaria if they develop ID or IDA [11,12].

Declining cord clamping for three minutes will prevent iron deficiency in the newborn as it maximizes restoration of red blood cells [13]. Pregnant women develop functional iron deficiency in the second trimester due to expansion of the plasma and red cell mass [14]. Treating iron deficiency in pregnancy improve the quality of life and prevents low birth weight [15]. Functional iron deficiency develops in athletes due to increased hepcidin. Patients with chronic diseases and inflammation can develop functional iron deficiency which is obscured by a normal to high ferritin as it is a positive inflammatory marker. Transferrin saturation of less than 20% and ferritin less than 30 mg is consistent with iron deficiency anemia. Other supporting features include increased soluble transferrin receptor, low hemoglobin, increased mean cell volume width and hypochromia, transferrin saturation of less than 20% with normal or increased ferritin, normal hemoglobin, normal soluble transferrin receptor and detectable stained iron in the bone marrow is consistent with functional iron deficiency, transferrin saturation of less than 20% with ferritin less than 70 ug/l, reduced MCV, normal or increased soluble transferrin receptor, reticulocytopenia and absent stainable iron in the marrow [16].

ID is characterized by normal hemoglobin, mean corpuscular volume, mean cell hemoglobin concentration, normal reticulocyte number and function, and normal soluble transferrin receptor. Reduced MCV, ferritin, transferrin saturation, reticulocytes, increased soluble transferrin receptor, elevated hepcidin and absent bone marrow stainable iron [17] are consistent with iron deficiency anaemia and functional iron deficiency. Parentage of hypochromic blood cells are equivalent to HbA1c in diabetic patients as it reflects restricted erythropoiesis in the preceding 2-3 months [18]. The number of reticulocytes reflects the iron availability for erythropoiesis in the preceding four days.

Oral iron causes few side effects and makes adherence to medication difficult. Ferrous sulphate causes constipation, diarrhoea, gastritis and black stools. An isotope study has shown frequent doses of iron during the day impairs absorption in the subsequent days and confirms that on alternate days there are higher rates of absorption [19]. Parenteral iron is safe and has become the standard of care. Iron should not be given during active sepsis as it can promote bacterial growth. Ferric carboxymaltose is very popular in Australia and is well-tolerated. A common dose is 15-20 mg/kg diluted in 250 ml of saline via a fifteen-minute infusion. An uncommon side effect is hypophosphatemia due to increased FGF23 which acts on the renal tubules, causing phosphaturia and hypophosphatemia. This does not cause problems and does not need phosphate replacement, except in frequent transfusion as it can cause osteomalacia and fractures [20,21].

Patients with iron deficiency should be investigated for their cause in order for it to be treated. If blood loss has been ruled out via endoscopy, capsule endoscopy, push endoscopy, and coeliac disease serology is negative, the patient should be referred to hematologist.

Conclusion

Iron deficiency and iron deficiency anemia can cause impaired quality of life and an increased disease burden across the globe. The safety of parenteral iron transfusion has transformed the treatment of iron deficiency, functional iron deficiency and iron deficiency anemia, and has been proven to be safe and efficient.

Acknowledgment: Associate Professor Adel Ekladios thanks Cassandra Byers for editing the manuscript.

References

1. WHO global Nutrition target 2025: policy brief series, 2014.
2. Andrewa Nc Disorders of iron metabolism, N Eng J Med. 1999; 341: 1986-95.
3. WHO. the Global prevalence of anemia in 2011.2015.
4. Mei Z, Cogswell ME, Looker AC, et al. Assessment of iron status in US pregnant women from the national health and nutrition examination survey (NHANES)1999-2006. Am J Clin Nutr. 2011; 93: 1312-20.
5. Kiss JE, Birch RJ, Steele WR, Wright DI, Cable RG, Quantification of body iron and iron absorption in the REDS-11 donor iron status evaluation (RISE) study. Transfusion. 2017; 57: 1656-64.
6. Muckenthaler MU, Rivella S, Hentze MW, Galy B. A red Carpet for iron metabolism, Cell. 2017; 168: 344-61.
7. Guide C. Altamura S, Klein FA et al. A novel inflammatory pathway mediating rapid hepcidin- independent Hypoferremia. Blood. 2015; 2265-75.
8. Mast AE. Low Hemoglobin deferral in Blood donors. Transfuse Med Rev. 2014; 28: 18-22.
9. Lombardi-Boccia G., Martinez-dominguez., Aguzzi A. total heme and non -heme iron in raw . and cooked meat J. Food Sci. 2002l; 67: 1738-1741.
10. Spottiswoode N, fried M, Darkesmith H, Drakmesith H, Duffy PE. Implications of Malaria on iron deficiency control strategies.

-
- Adv, Nute. 2012; 3: 570-578.
11. Armi mononuclear stage AE, Pinches R, Eddowes LA, Newbold associated with CI, darksmith H, Plasmodium Falciparum infected erythrocytes induced hepcidin (HAMP) mRNA Synthesis by peripheral blood mononuclear cells, Haematol. 2009 ; 147: 769-771.
 12. WHO . Guidelines : Delayed umbilical cord clamping for improving maternal and infant health and nutrition outcome, Geneva, World health organization. 2014.
 13. Rogozinska E, Daru J, Nicolaidis M et al Iron preparations for women of reproductive age with iron deficiency anemia in pregnancy (FRIDA); a systematic review and network meta-analysis Lancet Haematol. 2021; 8: e503_e512.
 14. Rukuni R, Knight M, Muephy MF, Roberts D, Stanworth SJ Screening for iron deficiency and iron deficiency anemia in pregnancy: a structured review and Gap analysis against UK national screening Criteria BMC pregnancy Childbirth. 2015; 15: 269.
 15. Pasricha S-R, Tye-Din, Muckenthaler MU, Swinkles DW Iron deficiency, Lancet. 2021; 397: 233-248.
 16. Thomas L, Transferrin saturation, Thomas L eds. Clinical laboratory diagnostics. 1998; 275-277.
 17. Baynes RD, Assessment of iron status. Clin Biochem. 1996; 29: 209-215.
 18. Cook JD, BS, Baynes RD. serum transferrin receptor, Annu Rev Med. 1993; 44: 63-47.
 19. Camaschella C. iron deficiency anaemia. N Engl J Med. 2015; 372: 1832-1843.
 20. Avni T, Bieber A, Grossman A, Green H, Leibovici L, Gafter-Gvili A. The safety of intravenous iron preparation's: Systemic review and meta-analysis. Mayo Clinic Proc. 2015; 90: 12-23.